REPORT DOCUMENTATION PAGE

AFRL-SR-BL-TR-99-

Public reporting burden for this collection of information is estimated to average 1 hour per response, in gathering and maintaining the data needed, and completing and revisiving the collection of information collection of information, including suggestions for reducing this burden, to Washington Headquarters 5 Davis Highway, Suite 1204, Artington, VA, 22202-4302, and to the Office of Management and Budget,

0030

a sources, ect of this Jefferson

Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget,	3.			
	d VIII 1118 1118			
	01 JUL 95 TO 30 JUN 98 FINAL			
4. TITLE AND SUBTITLE	5. FUNDING NUMBERS			
AASERT 95 (BMDO) CASCADED SPATIAL SOLITIONS FOR NETV				
ROUTING	3484/YS			
6. AUTHOR(S)				
Professor Stegeman				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZAT REPORT NUMBER				
University of Central Florida REPORT NUMBER				
4000 Central Florida Blvd Adm 243				
PO Box 25000				
Orlando FL 32816-0150				
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)	10. SPONSORING/MONITORING AGENCY REPORT NUMBER			
AFOSR/NE	AGENOT HEI OH HOME			
801 North Randolph Street Rm 732	F49620-95-1-0449			
Arlington, VA 22203-1977				
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION AVAILABILITY STATEMENT	12b. DISTRIBUTION CODE			
APPROVAL FOR PUBLIC RELEASED; DISTRIBUTION UNLIMITE				
MINOVALION I OBDIO NEEDIO NEEDIO NO INCIDENTIALI	- 1			
	19990209 085			
	17770207 007			
13. ABSTRACT (Maximum 200 words)				
Evnonimenta Lione performed as an additional in				
Experiments were performed on quadratic solito	n generation in Type I,			
non-critically phase-matched KNbO3. Using a very sensitive detector array and image enhancement protocols, the first picture of a quadratic soliton on				
and mage emidicement protocots, the first pic	ture of a quadratic soliton on			
was obtained. Further experiments were performed on KTP with the goal of				
demonstrating the spontaneous generation of quadratic solitons via down conversion from noise.				
Conversion from noise.				
l .				


14. SUBJECT TERMS			15. NUMBER OF PAGES
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
UNCLASSIFIED	UNCLASSIFIED	UNCLASSIFIED	UL
Standard Form 298 (Rey. 2-89) (EG)			

Project Title: Cascaded Spatial Solitons: Network Routing

The student supported by this AASERT, Russell Fuerst, has made considerable progress this year towards a PhD. His course requirements are finished and his Dissertation Proposal was accepted for a PhD program. His work was upgraded to an invited paper at IQEC'98 and OSA Annual Meeting. He attended a NATO Summer School in Corsica France in August of last. He is expected to graduate no later than the spring semester of 1999.

The ultimate goal of this program is to use spatial solitons as reconfigurable interconnects for guiding signals between multiple input and output ports. The immediate goal was to understand some of the basic properties of spatial solitons in quadratic media. The concept is to use steerable spatial solitons for guiding signal beams for implementation to NxN reconfigurable interconnects. His research progress in the last 10 months can be summarized as follows:

(1) Preliminary experiments were performed on quadratic soliton generation in Type I, non-critically phase-matched KNbO₃. Because the effective nonlinearity is 17 pm/V, a proliferation of solitons was observed at input intensities of 100s of MW/cm². Using a very sensitive detector array and image enhancement protocols, Russell was able to get the first picture of a quadratic soliton,



shown on the left. Light is input on the right hand side and the non-diffracting nature of the beam is clear for the sample which was 5 diffraction lengths long. However, the bandwidth of the OPO/OPG system is too broad for investigation of soliton properties. The OPO has been rebuilt to narrow the linewidth and experiments will continue with this material.

(2) Further experiments were performed on KTP with the goal of demonstrating the spontaneous generation of quadratic solitons via down conversion from noise. A strong input excitation at 532 nm was used, and the crystal was tilted away from the orientation needed for efficient second harmonic generation with a 1064 nm input. This tilt changes the phase-matching condition from degeneracy to non-degeneracy for the frequencies generated by down conversion. As a result, the generated quadratic soliton contained three different frequencies, the 532 nm component and two components equally split about 1064 nm by about 10 nm. Since this is a Type II crystal, the two fundamental components are also orthogonally polarized. At the time we did this experiment we believed that it was the first example of a quadratic soliton generated from noise in a single pass OPG/OPA configuration *Unfortunately*, a similar experiment was just published in Phys. Rev. Lett by Trapani et al.